

## **Certificate of Analysis**



This certificate is valid for two years from the date of shipment provided the substance is stored under the recommended conditions unopened in the original container.





The identity of the reference substance was established by following analyses.

## <sup>1</sup>H-NMR Spectrum la.

Conditions: 400 MHz, DMSO-d<sub>6</sub>



The structure is confirmed by the signals of the spectrum and their interpretation.





Method: 4.5 kV ESI+; vaporization temperature: 200 °C



The signals of the IR spectrum and their interpretation are consistent with the structural formula.

LOGICal® produced by LGC Page 3/6



## IIa. High Performance Liquid Chromatography (HPLC)

The purity of the reference substance was analysed by high performance liquid chromatography (HPLC).







## Area Percent Report - Sorted by Signal

Pk #	Retention Time	Area	Area %
1	3.25	15959	0.20
2	5.15	11410	0.14
3	5.91	8014489	98.88
4	6.61	62847	0.78
5	14.25	763	0.01
Totals		8105468	100.00
		~	

For the calculation the system peaks were ignored. The content of the analyte was determined as ratio of the peak area of the analyte and the cumulative areas of the purities, added up to 100 %.

(

	Ň	$ \setminus V \setminus \setminus \setminus$				
Results:						
Average	98.88 %					
Number of results	n=3					
Standard deviation	< 0.01 %	$\searrow$				
		$\mathbf{X}$				
		$\searrow$				
		)				
IIb. Water Content		)				
Method: Karl Fischer titration						
Desultar						
Results:						
Average	0.07 %					
Number of results	n=3					
Standard deviation 0.01 %						
	$\sim$					
~	//					
IIc. Residual Solvents	5					

Method: <sup>1</sup>H-NMR

No significant amounts of residual solvents were detected (< 0.05 %).





Chromatographic purity (HPLC)98.88 %Water content0.07 %Residual solventsNo significant amounts of residual solvents were detected (< 0.05 %).</th>Assay (100 % method)198.81 %

The assay is assessed to be 98.8 % 'as is'

The assay 'as is' is equivalent to the assay based on the not an hydrous and not dried substance respectively.

Release Date: Luckenwalde, 2015-03-05

> Dr. Sabine Schröder Product Release

<sup>1</sup> The calculation of the 100 % method follows the formula:

Assay (%) = (100 % - volatile contents) \*

Purity (%) 100 %

Volatile contents are considered as absolute contributions, purity is considered as relative contribution.



LGCFOR1387.02 lot number 72240 LGC GmbH, Im Biotechnologiepark, TGZ II, D-14943 Luckenwalde, Germany

 $\circledast$  2012 LGC limited. All rights reserved. LGC Standards is part of the LGC Group. LoGiCal is a registered trademark of LGC Standards GmbH