

Certificate of Analysis

Reference Substance

Apomorphine Hydrochloride Hemihydrate


Catalogue Number: LGCFOR0562.00

Lot Number: 1602

Molecular Formula: C₁₇H₁₇NO₂ HCl ½H₂O

Molecular Weight: 312.80 CAS Number: [41372-20-7] Long-term Storage: 2 to 8 °C, dark

Appearance: grey solid
Assay 'as is': 98.6 %



Date of shipment: 2016-May-20

This certificate is valid for two years from the date of shipment provided the substance is stored under the recommended conditions.

Release Date: 2010-02-19

LGC GmbH

Dr. Sabine Schröder Product Release

LGC Quality | ISO 9001:2008 DQS 102448 QM08






I. Identity

The identity of the reference substance was established by following analyses.

Ia. ¹H-NMR Spectrum

Conditions: 400 MHz, DMSO-d₆

The structure is confirmed with the signals of the spectrum and their interpretation.








lb. **IR Spectrum**

Method: Attenuated Total Reflection Fourier Transform Infrared (ATR-FTIR) Spectroscopy



Result of Peak Picking No. Position Intensity 3159 79 58 2282 2591.86 1503.24 1270.86 1157.08

The signals of the IR spectrum and their interpretation are consistent with the structural formula.

Purity II.

The purity of the reference substance was analysed by high performance liquid chromatography (HPLC).

HPLC Conditions:

Column: Conditions: Injector: **Detector:** Hypersil Gold (C18) 1.0 ml/min, 40 °C DAD Auto

5 μm, 150 x 4.6 mm 0-15 min Water/Acetonitrile 90/10 270 nm

30-35 min Water/Acetonitrile 90/10 (v/v);

5 µl; 0.0582 mg/ml in 15-20 min Water/Acetonitrile to 60/40 Water; 0.1 % H₃PO₄


20-25 min Water/Acetonitrile 60/40 25-30 min Water/Acetonitrile to 90/10

0.1 % H₃PO₄



LGCFOR0562.00 Lot Number 1602





Area Percent Report - Sorted by Signal

Pk#	Retention Time	Area	Area %	
1	3.07	244	0.01	
2	3.57	1053	0.02	
3	3.80//	292	0.01	
4	4.03	351	0.01	
5	4.7.1	12185	0.29	
6	5.76	470	0.01	
7	7.42	449	0.01	
8	8.15	4210130	99.56	
9	9.30	3138	0.07	
10	24.55	585	0.01	
Totals		4228897	100.00	

For the calculation the system peaks were ignored. The content of the analyte was determined as ratio of the peak area of the analyte and the cumulative areas of the purities, added up to 100 %.

Results:

Average 99.56 %

Number of results n=6

Standard deviation 0.04 %



LGCFOR0562.00 Lot Number 1602



III. Water Content

Method: Karl Fischer titration

Determined value	3.73 %	
Theoretical value	– 2.88 %	<
Content of excessive water	0.85 %	

IV. Residual Solvents

Method: 1H-NMR

Result: 0.07 % Ethanol

V. Final Result

Total impurities (HPLC)0.44%Water content0.85%Residual solvents0.07%Assay (100 % method) 1 98.64%

The assay is assessed to be 98.6 % 'as is'

The assay 'as is' is equivalent to the assay based on the not anhydrous and not dried substance respectively.

¹ The calculation of the 100 % method follows the formula:

Assay (%) = (100 % - KF - RES) * $\frac{Purity HPLC (\%)}{100 \%}$

Water (KF) and Residual solvents (RES) are considered as absolute contributions, HPLC purity is considered as relative contribution.

LGCFOR0562.00 Lot Number 1602